

Stream, and having no harbor that vessels drawing more than seven feet of water can enter. We stopped here to meet Mr. Charles Washburn, one of my collectors, who had been staying on the Islands for about three weeks. Very few resident species were obtained, but it is more than probable that a number of Bahama forms occur there which were not observed by us, as the time spent on these Islands was much too short to make anything like a thorough investigation.

Sterna maxima Bodd.	Vireo crassirostris Bryant.
Pelecanus fuscus Linn.	Vireo altiloquus barbatulus (Cab.).
Fregata aquila (Linn.).	Cereba bahamensis (Reich.).
Ardea herodias Linn.	Mniotilla varia (Linn.).
Ardea virescens bahamensis (Brewster).	Helinaia swainsoni Aud.
Ægialitis vocifera (Linn.).	Compsothlypis americana (Linn.).
Columbigallina passerina (Linn.).	Dendroica tigrina (Gmel.).
Columba leucocephala Linn.	Dendroica coronata (Linn.).
Crotophaga ani Linn.	Dendroica dominica (Linn.).
Ceryle alcyon (Linn.).	Dendroica palmarum (Gmel.).
Sphyrapicus varius (Linn.).	Dendroica discolor (Vieill.).
Doricha evelynæ (Bourc.).	Geothlypis trichas (Linn.).
Tyrannus dominicensis (Gmel.).	Setophaga ruticilla (Linn.).
Agelaius phœniceus bryanti Ridgw.	Seiurus motacilla (Vieill.).
Euetheia bicolor (Linn.).	Seiurus aurocapillus (Linn.).
Ammodramus savannarum passerinus (Wils.).	Polioptila cœrulea (Linn.).
	Galeoscoptes carolinensis (Linn.).
	Mimus polyglottos orpheus (Linn.).

LIST OF THE BIRDS COLLECTED BY C. L. WINCH IN THE CAICOS ISLANDS AND INAGUA, BA- HAMAS, DURING JANUARY AND FEBRUARY, AND IN ABACO, IN MARCH, 1891.

BY CHARLES B. CORY.

CAICOS ISLANDS.

Dafila bahamensis (Linn.). — Not uncommon; breeds on the Caicos Islands.

Aythya affinis (Eyton).

Ardea rufescens Bodd.—Several specimens of both the white and colored plumage of this species.

Ardea tricolor ruficollis (*Gosse*).
Symphechia semipalmata (*Gmel.*).
Hæmatopus palliatus (*Temm.*).
Columba gallina passerina (*Linn.*).
Columba leucocephala *Linn.*.
Zenaida zena (*Linn.*).
Pandion haliaëtus carolinensis (*Gmel.*).
Coccyzus minor maynardi *Ridgw.*
Crotophaga ani *Linn.*.
Sphyrapicus varius (*Linn.*).
Doricha evelynæ (*Bourc.*).
Tyrannus magnirostris *D'Orb.*—Three specimens taken. It has not previously been recorded north of Inagua.

Loxigilla violacea (*Linn.*).
Euetheia bicolor (*Linn.*).
Vireo crassirostris (*Bryant*).
Cæreba bahamensis (*Reich.*).
Compsothlypis americana (*Linn.*).
Dendroica petechia gundlachi *Baird*.
Dendroica coronata (*Linn.*).

Dendroica kirtlandii *Baird*—Two specimens taken. I believe it has no previous record so far south. The naturalists of the 'Albatros' expedition found it not uncommon in Rum Cay.

Dendroica palmarum (*Gmel.*).
Dendroica discolor (*Vieill.*).
Seiurus aurocapillus (*Linn.*).
Polioptila cærulea (*Linn.*).
Mimus gundlachi *Cab.*
Margarops fuscatus (*Vieill.*).

INAGUA.

Ardea herodias *Linn.*.
Sphyrapicus varius (*Linn.*).
Doricha lyrura *Gould*.
Tyrannus magnirostris *D'Orb.*
Myiarchus sagræ *Gundl.*

Loxigilla violacea (*Linn.*).—Mr. Winch writes me he believes the adult female of this species assumes a black plumage similar to the male. A number of black specimens taken by him proved on dissection to be females.

Euetheia bicolor (*Linn.*).
Vireo crassirostris (*Bryant*).
Cæreba bahamensis (*Reich.*).—Birds from Inagua have larger bills than those which occur on the northern Bahama Islands.
Dendroica tigrina (*Gmel.*).
Dendroica petechia gundlachi *Baird*.
Dendroica coronata (*Linn.*).

Dendroica dominica (*Linn.*).
Dendroica discolor (*Vieill.*).
Seiurus aurocapillus (*Linn.*).
Polioptila cærulea (*Linn.*).
Mimus gundlachi *Cab.*

Mimus polyglottos elegans (*Sharpe*). — Since writing the paper on *Mimus polyglottos* and *M. orpheus* which appeared in the January 'Auk' (p. 45) I have received some fifty specimens of the small *Mimus* from Inagua. With this additional material for comparison I find that a large majority of the Inagua specimens are somewhat smaller than *orpheus* or *polyglottos*, and have the primary coverts nearly and often completely covering the white on the quills. These differences are not constant but are probably sufficient to justify subspecific separation.

Margarops fuscatus (*Vieill.*).

A box of birds lately received from Inagua contained two species not observed by Mr. Winch. These are:—

Dendroica striata (*Forst.*). — Six specimens, April 23 to May 2.

Dendroica cærulescens (*Gmel.*). — Nine specimens, April 22 to May 1.

ABACO.

<i>Ægialitis semipalmata</i> <i>Bonap.</i>	<i>Cœreba bahamensis</i> (<i>Reich</i>).
<i>Columbigallina passerina</i> (<i>Linn.</i>).	<i>Mniotilla varia</i> (<i>Linn.</i>).
<i>Cathartes aura</i> (<i>Linn.</i>).	<i>Dendroica kirtlandi</i> <i>Baird.</i>
<i>Crotophaga ani</i> <i>Linn.</i>	<i>Dendroica petechia</i> <i>gundlachi</i> <i>Baird</i>
<i>Ceryle alcyon</i> (<i>Linn.</i>).	<i>Dendroica discolor</i> (<i>Vieill.</i>).
<i>Dryobates villosus</i> <i>maynardi</i> <i>Ridgw.</i>	<i>Dendroica coronata</i> (<i>Linn.</i>).
<i>Doricha evelynæ</i> (<i>Bourc.</i>).	<i>Dendroica dominica</i> (<i>Linn.</i>).
<i>Sporadinus ricordi</i> (<i>Gerv.</i>).	<i>Geothlypis trichas</i> (<i>Linn.</i>).
<i>Myiarchus sagræ</i> <i>Gundl.</i>	<i>Geothlypis tanneri</i> <i>Ridgw.</i>
<i>Pitangus bahamensis</i> <i>Bryant.</i>	<i>Seiurus noveboracensis</i> (<i>Gmel.</i>).
<i>Loxigilla violacea</i> (<i>Linn.</i>).	<i>Seiurus aurocapillus</i> (<i>Linn.</i>).
<i>Euetheia bicolor</i> (<i>Linn.</i>).	<i>Polioptila cærulea</i> <i>Linn.</i>
<i>Spindalis zena townsendi</i> <i>Ridgw.</i>	<i>Galeoscoptes carolinensis</i> (<i>Linn.</i>).
<i>Vireo crassirostris</i> (<i>Bryant.</i>).	<i>Mimus polyglottos orpheus</i> (<i>L.</i>).
<i>Vireo altiloquus barbatulus</i> (<i>Cab.</i>).	<i>Mimocichla plumbea</i> (<i>Linn.</i>).